


RULIDE



Rulide is an effective drug used for the treatment of urinary, respiratory tract and soft tissue infections.

DRUG USES

Rulide treats respiratory tract, soft tissue and urinary infections.

HOW TAKEN

Follow your doctor's prescription while taking the drug. The drug should be taken with food to avoid any stomach upset. Rulide should be swallowed whole and it should not be crushed, broken or chewed. Grape fruit and grape fruit juice should be avoided while taking Rulide. For the best results and complete cure of the infection, the whole course of the drug has to be taken. Even if you feel better, do not stop in the middle of a course.

WARNINGS/PRECAUTIONS

You have to inform your doctor about all your medical conditions before he prescribes the drug for you. You should have to tell your doctor if you are allergic to any particular medicine, food or to any substance.

Do not hesitate to tell your doctor if you kidney or liver problems, stomach or intestinal infections, heart problems, porphyria or myasthenia gravis.

The drug should only be taken with caution by pregnant women and breast feeding mothers. It is better to discuss with your doctor about all the complications related to Rulide before you take the drug during the pregnancy period. As Rulide is found in breast milk, you have to be cautious while taking the medicine during breast feeding.

You should also tell your doctor about the various medications that you are taking as it is seen that some medicines interact with Rulide. If you experience diarrhea, stomach pain or bloody stools, better contact your doctor.

It has also been seen that prolonged use of Rulide can cause a second infection. If signs of a second attack are seen, immediately contact your doctor so that he can make changes in the prescription.

The drug can interfere with lab tests, especially those connected with liver functions, kidney functions and blood cell counts. Therefore, it is better to tell the lab technician and the doctor that you are on Rulide medication before any tests are done.

MISSED DOSE

There is nothing to worry if a dose is missed. Take it as soon as you remember it. And if it is time for the next dosage, just skip the missed one and continue with your scheduled dosages. Never mix two does.